#### **Targets Of Opportunity**

Alan E. Pihlak, Leon A. Benjamin, Bobby R. Kelley

Forecast Systems Laboratory/OAR /NOAA/DoC

Daniel E. Wolfe, James R. Jordan, F. Martin Ralph

Environmental Technology Laboratory/OAR/NOAA/DoC

Kenneth S. Gage

Aeronomy Laboratory/OAR/NOAA/DoC

Brad Colman

Seattle, WA WFO/NWS/NOAA/DoC

Neil A. Stuart

Wakefield, VA WFO/NWS/NOAA/DoC

Richard Hayton, Robert Esser

Air Combat Command/USAF/DoD






## Cooperative Agency Profilers History

- 1970's Aeronomy Laboratory builds first wind profilers for research
- 1980's Colorado Wind Profiler Network built for research
- 1985 NOAA Profiler Network initiated
- 1994 NOAA Profiler Network assessment completed
- 1994 Boundary Layer Profiler (BLP) Acquisition Project started
- 1997 ~ 20 CAP active
- 1999 ~ 30 CAP active
- 2001 ~ 40 CAP active
- 2002 ~ 60 CAP active. Project name changed from BLP to Cooperative Agency Profiler – profilers 'not just for the Boundary Layer anymore' as more tropospheric and lower tropospheric profilers are deployed by agencies.





#### Number of CAP



Increase in number of CAP from 1995 to 2002





## Cooperative Agency Profilers Who Cooperates?

- NOAA Environmental Technology Laboratory 13 profilers
- NOAA Aeronomy Laboratory 7
- Kennedy Space Center And Cape Canaveral Air Force Station 5
- Argonne National Laboratory ABLE/ARM Programs 3
- NCAR Sponsored By The FAA 3
- South Coast Air Quality Management District 3
- White Sands Missile Range (Meteorology Branch) 3
- Dugway Proving Ground (DPG) 2
- Environment Canada 2
- Department Of Interior Minerals Management Service 2
- North Carolina Department Of Environment & Natural Resources (DENR) & Division Of Air Quality - 2
- San Diego Air Pollution Control District 2
- San Joaquin Valley APCD 2
- Texas Natural Resources Conservation Commission 2
- Several others with one profiler







Map of the active continental CAP sites excluding Alaska





## Cooperative Agency Profilers Agencies Primary Missions

- Environmental protection and air quality monitoring (many)
- Space and Missile launch (Vandenberg, Cape Kennedy and White Sands)
- Research experiments (ETL, AL, Air Resources Lab)
- Dept. of Defense (Dugway)
- Aviation (Juneau, AK)



915 MHz profiler with RASS at Las Vegas Courtesy of NOAA's Air Resources Laboratory





## Cooperative Agency Profilers Typical Characteristics

- Produce a wind profile every 5 to 30 minutes high temporal frequency compared to every 12 hours for radiosonde.
- For all profilers, height coverage is dependent on atmospheric turbulence, moisture, transmitter power and profiler frequency.
- Range gate spacing and other operating parameters are variable and set dynamically by the operator of the radar for their specific mission





#### **Typical Characteristics**

Frequency	915 MHz	449 MHz	50 MHz
Height AGL	>3 km	>8 km	>20 km
Peak Power	500 W	>2 kW	> 15 kW





Rutgers, NJ Profiler



915 MHz profiler with RASSCourtesy of Rutgers University





**Targets of Opportunity** 

- This project is an extension of the 1995 NPN processing infrastructure.
- Participation in the CAP project is strictly voluntary.
- No employees have ever worked full-time on this project. A level of 60 CAP in operation <u>almost</u> requires the attention of an operator full time.
- Total cost estimated at \$439,000 to process these data from 1994 to present – a cost of approximately \$1800/profiler/year





#### **Processing Infrastructure**



#### Processing

Convert time to UTC
Convert to common file format
Add QC

#### Acquisition

FTP GOES WWW Dial-up

#### Data Delivery

NWS NCEP FSL (Backup RUC, MADIS) ETL UCAR

Dugway Proving Grounds
National Atmospheric Release Advisory Center (LLNL)
Chesapeake Bay LAPS Model
www.profiler.noaa.gov





## Cooperative Agency Profilers Benefits to Agencies

- NPN adds advanced quality assessment information.
- Wind and RASS data available to the public, as graphics and raw data, on NPN website.
- Agencies often have limited budget for operations the NPN provides monitoring and notification of profiler failure.
- Radar wind profiler experts in the Profiler Control Center can identify difficult technical problems and suggest solutions.





## Cooperative Agency Profilers Benefits to Public

High temporal frequency wind and RASS data from otherwise data-sparse regions aids in :

- Improved Forecasts (National Weather Service)
  - Wind shear
  - Low level jets
  - Advection patterns
  - Inversion depth
  - Temperature profiles (RASS)
- Numerical Weather Prediction (NWP) CAP data is 'in the tanks' at National Centers for Environmental Prediction - evaluation expected to begin in February 2003.
- National security NWP models of atmospheric 'release' events (NARAC).
- General public private aviation, wind surfing, hot-air ballooning, hang gliding, ultralight airplanes.






### Cooperative Agency Profilers The Future

- Ability to display CAP data on NWS Advanced Weather Processing Information System (AWIPS)
- Provide GOES data collection implementation for CAP profilers.
- Improved quality control algorithms such as coherent interference detection, precipitation contamination.
- Automated comparison of profiler data vs. models and sondes for quality assessment.





Location of TARS profilers



## Cooperative Agency Profilers The Future




TARS Aerostat – Wind profiler needed for operations





## Cooperative Agency Profilers The Future



WINDAS - Japanese Meteorological Agency Profiler Network





#### Summary

- Numbers of CAP are increasing NPN set to add several more in the next few weeks.
- NPN has the infrastructure in place to acquire and distribute these data.
- Essentially zero impact on the Cooperative agency's primary mission and budget.
- The cost is low (~ \$1800/profiler/year).
- Benefits the taxpayer primarily by improved forecasting with additional effect via improved data access for modeling and national security.
- Benefits the Cooperative agencies in that they do not have to become or hire profiler experts to monitor their systems and diagnose problems.
- Data archive available to public on http://www.profiler.noaa.gov



